

AD _____

Award Number: DAMD17-02-1-0662

TITLE: Dietary Genistein and Prostate Cancer Chemoprevention

PRINCIPAL INVESTIGATOR: Coral A. Lamartiniere, Ph.D.

CONTRACTING ORGANIZATION: The University of Alabama at Birmingham
Birmingham, AL 35294-0111

REPORT DATE: April 2004

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

20040608 137

REPORT DOCUMENTATION PAGEForm Approved
OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE April 2004	3. REPORT TYPE AND DATES COVERED Annual (1 Apr 2003 - 31 Mar 2004)	
4. TITLE AND SUBTITLE Dietary Genistein and Prostate Cancer Chemoprevention			5. FUNDING NUMBERS DAMD17-02-1-0662	
6. AUTHOR(S) Coral A. Lamartiniere, Ph.D.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) The University of Alabama at Birmingham Birmingham, AL 35294-0111 E-Mail: coral.lamartiniere@ccc.uab.edu			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012			10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited				12b. DISTRIBUTION CODE
13. ABSTRACT (Maximum 200 Words) The disease of cancer is usually attacked at time of diagnosis, and even chemoprevention is not usually considered until adulthood. Our hypothesis is that windows of development hold the key for chemoprevention of prostate cancer. We have previously demonstrated that genistein is bioavailable to the rat prostate and that life-time exposure to physiological concentrations of genistein suppressed the development of chemically-induced prostate cancer. The purpose of our research is to determine if there is a developmental window for this chemoprevention and the mechanism(s) of chemoprevention. We have demonstrated that neonatal and prepubertal exposure to genistein via the diet does not alter development of the prostate buds in 21 and 35 day old rats. More recently, we have demonstrated that prepubertal, as well as adult and lifetime, exposure to genistein in the diet down regulated androgen receptor, but not estrogen receptors alpha and beta, expression in the dorsolateral prostate of rats. Down regulated androgen receptor is consistent with genistein suppressing prostate cancer development. Short term exposure to genistein having a permanent effect on androgen receptor expression is hypothesized to occur via androgen receptor promoter DNA methylation. Tumorigenesis studies are completed and are in the process of being evaluated for histopathology.				
14. SUBJECT TERMS Genistein, Rat, Prostate, Cancer Chemoprevention				15. NUMBER OF PAGES 6
				16. PRICE CODE
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

NSN 7540-01-280-5500

Standard Form 298 (Rev. 2-89)
Prescribed by ANSI Std. Z39-18
298-102

Table of Contents

Cover.....	1
SF 298.....	2
Table of Contents	3
Introduction.....	4
Body.....	4
Key Research Accomplishments.....	6
Reportable Outcomes.....	6
Conclusions.....	6

Introduction

The disease of cancer is usually attacked at time of diagnosis, and even chemoprevention is not usually considered until adulthood. Our hypothesis is that windows of development hold the key for chemoprevention of prostate cancer. We have previously demonstrated that life-time exposure to physiological concentrations of genistein suppressed the development of chemically-induced prostate cancer. We have shown that genistein is bioavailable to the prostate. We have also demonstrated that genistein did not readily cross the placenta, hence we don't believe that the gestational period plays a significant role in the protective effect. On the other hand, it is primarily during the first weeks of postnatal life that prostate differentiation takes place, a period that may be influenced by genistein. The purpose of our proposed research is to determine if there is a developmental window for this chemoprevention and the mechanism (s) of chemoprevention. The importance of this lies in the need to know, prior to initiation of human trials, if we need to expose infants and/or adults to get maximum chemoprevention. We propose to accomplish this in a dietary model at "physiological concentrations".

Body

Aim 1. To determine if a specific window of development (prepubertal only, adult only or life-time) is responsible for genistein chemoprevention of prostate cancer. This will be done on the following groups of rats, 30/group initially.

Group A) genistein via the diet from birth throughout life to confirm that postnatal lifetime exposure only protects against prostate cancer.

Group B) genistein in the diet from birth until 35 days of age only.

Group C) genistein in the diet starting at 90 days of age, 20 days after cancer initiation.

Group D) no genistein in the diet as positive controls.

This was initiated in the middle of the first year. The breeders were purchased, the rats bred, treated with the carcinogenesis protocol, and exposed to genistein in the diet as listed above. The last of the rats for this aim were recently necropsied this month. The tissues have been fixed and the blocks are being sectioned. The slides will be submitted for hemotoxylin and eosin staining and sent to Dr. Eltoum for histopathologic evaluation. The histopathology report will be available in the beginning of year 3.

Aim 2. To investigate prostate gland morphology in the dorsal and lateral lobes of the prostates of 21 and 35 day old rats exposed \pm genistein in the diet, starting at birth. Months 6-12. No statistical significance was detected for prostate bud perimeter from genistein compared to control treated rats. This work was completed in year 1 and has already been reported.

Aim 3. To investigate the potential of genistein to regulate sex steroid receptor expression as mechanism of prostate cancer prevention. This was projected to be carried out during year 2.

Dietary genistein resulted in decreased androgen receptor, but not estrogen receptors alpha or beta, expression in dorsolateral prostates (Figures 1 and 2). The novel finding is that prepubertal short term (days 1-35 postnatal) genistein exposure was able to down regulate the androgen receptor as well as did the adult only (days 56-70) and life time (days 1-70) genistein treatments. Since short term early exposure to genistein is as effective as short term late and lifetime exposure to genistein, this suggests that early exposure to genistein can have a programming or imprinting effect on androgen receptor expression. One mechanism by which genistein could




Figure 1. Dietary genistein reduces androgen receptor expression in dorsolateral prostates of rats. a = $p < 0.01$ and b = $p < 0.001$ compared to controls, respectively as determined by one-way ANOVA. Each group contained 6 samples, 2 prostates/sample).




Figure 2. Estrogen receptor expression in dorsolateral prostates of rats fed genistein in the diet. NS: No significant effect compared to controls as determined by one-way ANOVA. Each group contained 6 samples, 2 prostates/sample).

cause a permanent effect on gene and protein expressions is via DNA methylation (See next Aim).

Aim 4. To investigate DNA methylation as imprinting mechanism of action. This is to be carried out during year 3. From the data in Figures 1 and 2, it is evident that the “permanent” down regulated expression of the androgen receptor could be explained on the basis of DNA methylation. In year 3, we will treat rats as described above with genistein, isolate DNA from the prostates, subject the DNA to bisulfite modification followed by PCR with methylation specific primers and gel analysis for methylation specific PCR analysis.

Key Research Accomplishments

Prepubertal, as well as adult and lifetime, exposure to genistein in the diet down regulated androgen receptor, but not estrogen receptors alpha and beta, expression in the dorsolateral prostates of rats.

Reportable Outcomes

None.

Conclusion

Down regulated androgen receptor is consistent with genistein suppressing prostate cancer development. Short term exposure to genistein having a permanent effect on androgen receptor expression is hypothesized to occur via androgen receptor promoter DNA methylation.